



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

in re Application of

Atty. Docket

WALID ALI

US 010194

Serial No.: 09/938,377

Group Art Unit: 2661

Filed: AUGUST 24, 2001

Title: AN APPARATUS AND METHOD FOR COMBINING RANDOM SET OF VIDEO FEATURES IN A NON-LINEAR SCHEME TO BEST DESCRIBE PERCEPTUAL QUALITY OF VIDEO SEQUENCES USING HEURISTIC SEARCH METHODOLOGY

Commissioner for Patents
Washington, D.C. 20231

RECEIVED

MAY 02 2002


LETTER TO OFFICIAL DRAFTSMAN

Technology Center 2600

Sir:

Enclosed are (4) four sheets of formal drawings
for filing in the above-identified application.

Respectfully submitted,

By 
Russell Gross, Reg. No. 40,007
Attorney
(914) 333-9631

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being
deposited this date with the United States Postal Service as
first-class mail in an envelope addressed to:

COMMISSIONER OF PATENTS AND TRADEMARKS
Washington, D.C. 20231

On April 23, 2002

By John Chyz

1/4




FIG. 1A

2/4




FIG. 1B

WHEN A PREDETERMINED NUMBER OF SETS OF METRICS = n, THE QUADRATIC MODEL TO OBTAIN THE OBJECTIVE EVALUATION F IS:

$$F = \left(\sum_{i=1}^n w_i x_i \right)^2, \text{ WHEREIN "n" IS A NON-ZERO VALUE.}$$

FIG. 1C

WHEN A NUMBER OF THE SET OF METRICS = 4, THEN THE QUADRATIC MODEL TO OBTAIN THE OBJECTIVE EVALUATION F IS:

$$F = w_1^2 x_1^2 + w_2^2 x_2^2 + w_3^2 x_3^2 + w_4^2 x_4^2 + w_5^2 x_1^2 x_2^2 + w_6^2 x_1^2 x_3^2 + w_7^2 x_1^2 x_4^2 + w_8^2 x_2^2 x_3^2 + w_9^2 x_2^2 x_4^2 + w_{10}^2 x_3^2 x_4^2$$

FIG. 1D

3/4

190

SELECTING A BEST OF WEIGHTS FROM THE PLURALITY OF SETS OF WEIGHTS PROVIDED IN STEP (130), SAID BEST SET OF WEIGHTS BEING HEURISTICALLY DETERMINED BY A GENETIC ALGORITHM THAT INCREASES DYNAMICALLY A SIZE OF THE ASSIGNED RANGE OF SAID EACH ONE SET OF WEIGHTS PROVIDED IN STEP (130).

FIG. 1E

200

SELECTING A BEST OF WEIGHTS FROM THE PLURALITY OF SETS OF WEIGHTS PROVIDED IN STEP (130), SAID BEST SET OF WEIGHTS BEING HEURISTICALLY DETERMINED BY A GENETIC ALGORITHM THAT ENABLES FINDING THE BEST SOLUTION THAT MAXIMIZES THE CORRELATION FACTOR R OF THE OVERALL OBJECTIVE IMAGE QUALITY F WITH THE SUBJECTIVE EVALUATION WITHOUT THE NEED TO CARRY OUT AN EXHAUSTIVE SEARCH TO FIND THE BEST SET OF WEIGHTS.

FIG. 1F

CALCULATING OF THE CORRELATION FACTOR R IN STEP (160) BY USING A SPEARMAN RANK ORDER COMPRISING THE FOLLOWING EQUATION:

$$R = 1 - \frac{6 \cdot \sum (X_i - \bar{X})(Y_i - \bar{Y})}{k(k^2 - 1)}$$

WHEREIN X IS EQUAL TO A VECTOR OF RANKED k OBJECTIVE VALUES FOR THE k SEQUENCES (k*1), AND

Y IS EQUAL TO A VECTOR OF RANKED k SUBJECTIVE EVALUATION FOR THE k SEQUENCES (k*1).

FIG. 2




FIG. 3